

# VU Research Portal

## Central pain processing is altered in people with Achilles tendinopathy.

Tompra, N.; van Dieen, J.H.; Coppieters, M.W.J.

### ***published in***

British Journal of Sports Medicine  
2015

### ***DOI (link to publisher)***

[10.1136/bjsports-2015-095476](https://doi.org/10.1136/bjsports-2015-095476)

### [Link to publication in VU Research Portal](#)

### ***citation for published version (APA)***

Tompra, N., van Dieen, J. H., & Coppieters, M. W. J. (2015). Central pain processing is altered in people with Achilles tendinopathy. *British Journal of Sports Medicine*. <https://doi.org/10.1136/bjsports-2015-095476>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

# Central pain processing is altered in people with Achilles tendinopathy

Nefeli Tompra,<sup>1</sup> Jaap H van Dieën,<sup>1</sup> Michel W Coppieters<sup>1,2</sup>

<sup>1</sup>Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, MOVE Research Institute Amsterdam, VU University Amsterdam, Amsterdam, The Netherlands

<sup>2</sup>Department of Physiotherapy, School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane, Queensland, Australia

## Correspondence to

Professor Michel Coppieters, MOVE Research Institute Amsterdam, Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam, Van der Boerhorststraat 9, Amsterdam 1081BT, The Netherlands; [m.coppieters@vu.nl](mailto:m.coppieters@vu.nl)

Accepted 23 November 2015

## ABSTRACT

**Background** Tendinopathy is often a chronic condition. The mechanisms behind persistent tendon pain are not yet fully understood. It is unknown whether, similar to other persistent pain states, central pain mechanisms contribute to ongoing tendon pain.

**Aim** We investigated the presence of altered central pain processing in Achilles tendinopathy by assessing the conditioned pain modulation (CPM) effect in people with and without Achilles tendinopathy.

**Methods** 20 people with Achilles tendinopathy and 23 healthy volunteers participated in this cross-sectional study. CPM was assessed by the cold pressor test. The pressure pain threshold (PPT) was recorded over the Achilles tendon before and during immersion of the participant's hand into cold water. The CPM effect was quantified as the absolute difference in PPT before and during the cold pressor test.

**Results** An increase in PPT was observed in the Achilles tendinopathy and control group during the cold pressor test ( $p<0.001$ ). However, the CPM effect was stronger in the control group (mean difference=160.5 kPa, SD=84.9 kPa) compared to the Achilles tendinopathy group (mean difference=36.4 kPa, SD=68.1 kPa;  $p<0.001$ ).

**Summary** We report a reduced conditioned pain modulation effect in people with Achilles tendinopathy compared to people without Achilles tendinopathy. A reduced conditioned pain modulation effect reflects altered central pain processing which is believed to contribute to the persistence of pain in other conditions. Altered central pain processing may also be an important factor in persistent tendon pain that has traditionally been regarded to be dominated by peripheral mechanisms.

## INTRODUCTION

Tendinopathy is a generic term used to describe tendon pathologies that typically manifest in pain, swelling and dysfunction.<sup>1</sup> Despite best available management, tendon pain often becomes chronic.<sup>2–3</sup> Furthermore, the mechanisms underlying persistent tendon pain are not yet fully understood.

Tendinopathy has traditionally been viewed as a peripheral disorder confined to the tendon itself. However, recent evidence suggests other possible contributors to the persistence of tendon pain.<sup>4–8</sup> Many chronic pain conditions share a number of typical features, such as hyperalgesia/allodynia,<sup>9–10</sup> widespread pain<sup>11–12</sup> and reduced endogenous pain modulation.<sup>10–13</sup> In line with these presentations, various tendinopathies are characterised by mechanical hyperalgesia,<sup>14–15</sup> pinprick allodynia and increased vibration disappearance thresholds.<sup>16</sup> In

contrast, widespread pain is not a common clinical presentation of tendinopathy.<sup>5–17–18</sup> Yet, bilateral symptoms (including sensory and motor deficits) are often reported.<sup>3–6–14</sup> Up to now, altered central pain modulation (CPM) mechanisms have received little research attention in people with tendon pathology.

Endogenous pain modulation is governed by the central nervous system via facilitatory and inhibitory mechanisms. CPM testing paradigms are used to examine the efficacy of descending pain modulatory pathways.<sup>19</sup> In this experimental approach, the application of a noxious 'conditioning' stimulus typically results in pain inhibition for a subsequent testing stimulus. CPM is, therefore, quantified as the change in the intensity necessary to evoke pain (or pain intensity) of the testing stimulus before and during the application of the conditioning stimulus.<sup>20</sup> Persistent pain may reflect a possible dysfunction in the inhibitory circuits<sup>21</sup> which has been evident in a plethora of chronic pain conditions such as fibromyalgia,<sup>22</sup> irritable bowel syndrome,<sup>23</sup> pancreatitis,<sup>24</sup> tension type headache,<sup>25</sup> temporomandibular disorders,<sup>23</sup> osteoarthritis<sup>26</sup> and low back pain.<sup>27</sup> Whether chronic tendon pain is associated with similar deficits in central pain modulatory mechanisms is yet to be investigated. Therefore, the aim of this study was to examine the efficacy of CPM in people with chronic Achilles tendinopathy compared to healthy individuals.

## METHODS

### Study design

A cross-sectional study.

### Participants

People with and without Achilles tendinopathy were eligible to participate. Participants were recruited via print and social media advertisements from October 2014 to February 2015 they had to be actively engaged in running activities at the period of testing. All volunteers with pain over the Achilles tendon region underwent a patient interview and clinical examination. The diagnostic criteria for Achilles tendinopathy included a characteristic history of activity-related pain and tenderness on tendon palpation.<sup>15–28</sup> Participants with Achilles tendinopathy were included if they presented with pain in the Achilles tendon for at least 3 months prior to testing. Participants with any other medical condition or musculoskeletal disorder in the preceding 6 months that lasted for more than 1 week or for which treatment was sought were excluded. Further exclusion criteria were the presence of systemic disorders,

**To cite:** Tompra N, van Dieën JH, Coppieters MW. *Br J Sports Med* Published Online First: [please include Day Month Year] doi:10.1136/bjsports-2015-095476

cardiovascular or neurological problems, fibromyalgia and any medication usage.

Thirty volunteers with pain over the Achilles tendinopathy region agreed to participate, but 10 did not meet the criteria (9 were excluded because they did not have Achilles tendinopathy; 1 patient with Achilles tendinopathy experienced symptoms for less than 3 months). In addition, 26 symptom-free runners without Achilles tendinopathy were recruited from the same population, using the same recruitment strategies.

All participants completed the Victorian Institute of Sports Assessment—Achilles Questionnaire (VISA-A)<sup>29</sup> and the 21-item activities of daily living (ADL) subscale of the Foot and Ankle Ability Measure (FAAM).<sup>30</sup> Additionally, the participants from the Achilles tendinopathy group were asked to draw the areas where they experienced pain on a lower limb chart.<sup>28</sup>

The study was approved by the local ethics committee of the Department of Human Movement Sciences of VU University, Amsterdam, the Netherlands (Reference: ECB 2014–58). It was conducted according to the Declaration of Helsinki and all participants signed an informed consent prior to their participation.

### CPM testing protocol

To explore CPM, the cold pressor test was used as the conditioning stimulus and the pressure pain threshold (PPT) as the test stimulus.<sup>19 20</sup> Participants submerged their hand contralateral to their affected ankle (or the most affected in case of bilateral Achilles tendinopathy) in a container with cold water (approximately 9° Celsius). Healthy volunteers immersed their dominant hand. Continuous circulation of the water in the container was implemented in order to prevent an increase in temperature around the hand.<sup>31</sup> The water temperature was monitored by a thermometer with a digital display (resolution: 0.1°C). The participants placed their hand in the water up to the level of the wrist and they were instructed to keep their fingers spread. They were asked to rate their perceived hand pain on a 0 ('no pain') to 10 ('worst imaginable pain') numerical pain rating scale (NPRS). The CPM assessment began as soon as a score of 5 was reached. When required, ice was added to the water in order to maintain a minimum pain score of 4 on the NPRS during the experiment.

The PPT measurements were performed using a digital algometer (Type II, Somedic AB, Stockholm, Sweden). The participants were placed prone on a treatment plinth with their feet against the wall ensuring an ankle position of 0°. The assessment site was the most painful location of the Achilles tendon as determined by manual palpation.<sup>32</sup> For the control group, the point of pressure application was the mid-portion of the Achilles tendon (approximately 2–3 cm proximal to the insertion). The probe was placed perpendicular to the skin and the pressure was applied through a 1 cm<sup>2</sup> rubber plate at a constant rate of 40 kPa/s. The participants were instructed to press a button as soon as the applied pressure was perceived as painful.

Three consecutive trials with 30 s intervals were performed before and during the application of the conditioning stimulus. If the difference in consecutive PPTs reached more than 100 kPa, the procedure was repeated until three PPTs would differ by less than 100 kPa within a maximum of five measurements in total. The mean score of the three trials with the least variance was calculated and used for further analysis. This method showed excellent intra-rater reliability (Intraclass Correlation Coefficient (ICC<sub>(2,1)</sub>): 0.96; 95% CI 0.88 to 0.99; SE of Measurement (SEM): 8.74 kPa; Minimum Detectable

Change (MDC): 24.2 kPa) in a prior pilot study on 12 healthy volunteers.

To familiarise the participants with the PPT measurements, PPTs were performed over the thenar eminence of the dominant hand and over the Achilles tendon contralateral to the tested ankle before the actual experiment started. All experiments took place in the Vrije University of Amsterdam, Amsterdam, the Netherlands.

### Statistical analysis

The characteristics of the group with and without Achilles tendinopathy were compared with independent *t* tests, given a normal distribution of the data. Independent *t* tests were also used to examine potential differences between groups for the water temperature and the intensity of the induced hand pain during cold pressor test. For ordinal or non-normally distributed data, the equivalent non-parametric test (Mann–Whitney *U*) was performed. Normality was checked by visual inspection of the *q*–*q* plot, the box plot of the data, and a Shapiro–Wilks test.

To determine whether there was a difference in CPM responses between the Achilles tendinopathy and the control group, PPTs were investigated in a 2×2 (Time (before and during the cold pressor test)×Group (Achilles tendinopathy and control group)) repeated-measures analysis of variance (ANOVA). The normality and homogeneity of variance of the data were examined prior to the test. There were no violations of these assumptions. Any Group×Time interaction was clarified by examining the difference in the CPM effect (quantified as the difference in PPT values before and during cold pressor test) between the groups with an independent *t* test.

Given a non-normal distribution, Spearman correlation coefficients (*r*) were calculated to explore whether the pain severity (NPRS), symptom duration, functional limitation (FAAM) and Achilles tendinopathy severity (VISA-A) were related to the CPM response in the Achilles tendinopathy group.

### RESULTS

Three healthy participants were excluded from the analysis (two participants did not maintain the minimum required score of 4 on the NPRS during the cold pressor test; one participant reported no pain during pressure application over the Achilles tendon). **Figure 1** visualises the pain location as reported by the Achilles tendinopathy group. The characteristics of all participants who completed the experiment are shown in **table 1**. The Achilles tendinopathy group was on an average older than the control group (mean difference 6.8 years), yet the difference did not reach statistical significance (95% CI (–14.68 to 1.18)). The amount of running and cross-training per week was not significantly different between the two groups. The outcomes of FAAM and VISA-A were significantly lower in patients compared to healthy participants. The PPT on the thenar eminence did not differ significantly between the groups (mean difference=35.1, *t*(41)=1, *p*=0.33, 95% CI (–37.3 to 107.6)).

### Conditioned pain modulation

During the cold pressor test, the water temperature was similar for both the Achilles tendinopathy and control group (approximately 9°C). The level of induced hand pain was slightly higher for people with Achilles tendinopathy than for healthy participants, yet the difference was not statistically significant (mean difference=0.7, 95% CI (–1.5 to 0.2); **table 2**).

There was a significant main effect for time (*F*(1,41)=68.93, *p*<0.001,  $\eta^2_r$  = 0.63) and an interaction with the group (*F*(1,41)=27.36, *p*<0.001,  $\eta^2_r$  = 0.4), indicating that the

change in PPT during the cold pressor test was different for both groups. The CPM effect for the Achilles tendinopathy group was significantly lower than for the control group (mean difference=124.1 kPa,  $t(41)=5.2$ , 95% CI (76.2 to 192)). The CPM responses are illustrated in [table 2](#).

### Correlations


No significant correlations between pain severity experienced the week prior to the test (NPRS) ( $r=0.19$ ;  $p=0.427$ ), duration of symptoms ( $r=-0.41$ ;  $p=0.07$ ), FAAM score ( $r=0.22$ ;  $p=0.35$ ) or the VISA-A score ( $r=0.02$ ;  $p=0.95$ ) and the CPM effect were identified.

### DISCUSSION

We investigated differences in central pain processing between people with Achilles tendinopathy and healthy individuals. There was a significantly smaller CPM effect exhibited during the cold pressor test in the Achilles tendinopathy group compared to the control group. This indicates that central pain modulatory processes are altered in people with Achilles tendinopathy.

### CPM and chronic pain

CPM is a phenomenon that reflects the activity of the descending pain modulation system.<sup>20 33</sup> A deficit in its function has



**Figure 1** Area of pain for the people with Achilles tendinopathy. Darker colours represent a larger number of participants reporting pain the corresponding area. Regardless of whether the left or right Achilles tendon was affected, the locations are summarised on a diagram of the right ankle (ie, locations on the left side were mirrored).

**Table 1** Characteristics (mean (SD)) of the Achilles tendinopathy (Achilles tendinopathy) and the control group

	Control (N=23)	Achilles tendinopathy (N=20)	p Value
Age (years)	36.2 (12.3)	42.9 (13.5)	0.09
Gender (male/female)	16/7	16/4	–
Body mass index	21.9 (2.5)	23.4 (2.7)	0.11
Symptom duration (months)	NA	21.8 (26.1)	–
Side of symptoms (unilateral/bilateral)	NA	11/9	–
NPRS (past week)	NA	3.7 (1.9)	–
VISA-A questionnaire	99.8 (0.9)	71.6 (15.8)	<0.001
FAAM questionnaire	99.8 (0.5)	90.5 (10.7)	<0.001
Running (km/week)*	30.0 (33)	27.5 (45)	0.11
Cross-training (hours/week)	2.4 (2.6)	1.7 (1.7)	0.51
PPT thenar (kPa)	372.5 (107.7)	337.4 (127.5)	0.33

FAAM, Functional Ankle Ability Measure; Km, kilometres; NA, not applicable; NPRS, Numerical Pain Rating Scale; PPT, pain pressure threshold; VISA-A, Victorian Institute of Sport Assessment-Achilles.

\*Data are presented as Median (IQR).

been reported in a variety of clinical conditions associated with chronic pain.<sup>34–36</sup> The results of this study reveal a reduced CPM effect in people with Achilles tendinopathy. As no previous studies have investigated CPM in a tendinopathy, comparisons with the present study are not yet possible. Nonetheless, our findings provide evidence that people with Achilles tendinopathy share similar changes in central pain modulation mechanisms as do chronic pain patients with fibromyalgia,<sup>22 37</sup> irritable bowel syndrome,<sup>23 38 39</sup> temporomandibular disorders,<sup>23 40</sup> whiplash-associated disorders,<sup>41 42</sup> tension type headache,<sup>25 43 44</sup> osteoarthritis<sup>26 45</sup> and low back pain.<sup>27</sup> Such changes have been suggested to contribute to the development of central sensitisation<sup>9 46–48</sup> and chronicity of pain.<sup>49 50</sup> Indeed, most of the aforementioned conditions are expected to exhibit signs of central sensitisation.<sup>11 26 51–58</sup> Whether this is also the case in people with Achilles tendinopathy remains to be investigated in future research. Nonetheless, the reduced CPM effect in people with Achilles tendinopathy, as observed in this study, suggests a role of altered central pain mechanisms and may reveal a factor contributing to chronic tendon pain.

It is a novel and interesting finding that CPM is reduced in people with Achilles tendinopathy. Tendinopathy seems a particular chronic pain state in which tendon pain is provoked

**Table 2** Cold pressor test (mean (SD))

	Control (N=23)	Achilles tendinopathy (N=20)	p Values*
Achilles tendon PPT (kPa)†			
Before cold pressor test	671.4 (215.7)	253 (80.5)	<0.001
During cold pressor test	831.9 (213.3)	289.4 (114.3)	<0.001
CPM effect (kPa)	160.5 (84.9)	36.4 (68.1)	<0.001
Water temperature (°C)	9.0 (0.9)	9.1 (0.8)	0.73
NPRS during cold pressor test	6.2 (1.3)	6.9 (1.6)	0.15

\*p Values correspond to independent t tests.

†Contralateral to the dominant hand for the control group and the (most) affected for the Achilles tendinopathy group.

°C, degrees Celsius; CPM, conditioned pain modulation; kPa, kilopascal; NPRS, Numerical Pain Rating Scale.



## Original article

during loading and subsides once the load has been removed.<sup>5</sup> Moreover, unlike other chronic pain conditions, Achilles tendinopathy is not accompanied by great limitations in physical functioning or ADLs. Particularly in the recruited Achilles tendinopathy group, despite the long mean symptom duration ( $\approx 2$  years), pain intensity was moderate ( $\approx 4$  on the NPRS) and the activity level similar to the control group. In contrast to what has been reported so far in the literature,<sup>59</sup> in the present study the CPM effect was not correlated with the functional ability or the activity level of the patient group.

### Central versus local mechanisms

Of particular importance in interpreting the altered CPM response may be the net influences of local and central pain mechanisms. It has been argued that local neural and inflammatory processes may conceal endogenous analgesia.<sup>60 61</sup> Sustained peripheral nociceptive activity from the tendon tissue (see Rio *et al*<sup>5</sup> for a review) may sensitise nociceptive neurons in the dorsal horn of the spinal cord<sup>9</sup> increasing their excitability, thus counteracting the effects of descending inhibition. In two studies in people with knee and hip osteoarthritis, the CPM effect was weakened before but not after surgery, implying that the reduced CPM was the result of ongoing nociceptive activity.<sup>45 62</sup> Although inflammatory mediators are not always expressed in tendinopathy,<sup>5 63</sup> it is possible that the observed reduction in the CPM effect is the manifestation of both peripheral and central pain mechanisms rather than central changes alone.

Methodological variability with regards to CPM testing location has been noted in the literature. Attenuated CPM in conditions such as irritable bowel syndrome, tension type headache and pancreatitis has been demonstrated at pain-free sites,<sup>23–25 38</sup> whereas in temporomandibular disorders and osteoarthritis reductions were found only at the painful or intrasegmental areas.<sup>26 40 45</sup> The syllogism behind testing CPM at a pain-free or intact tissue is based on the fact that CPM is regarded as a systemic rather than a localised phenomenon.<sup>60</sup> Hence, a CPM reduction exhibited in a pain-free area would be indicative of a limitation of an individual's endogenous analgesic capacity. Moreover, as speculated above, a reduction in CPM in the affected tissue could be conveyed by both peripheral and central pain mechanisms. On the other hand, the absence of a CPM reduction in an area remote to the painful site does not exclude a possible contribution of central pain modulatory processes to the ongoing pain. The idea that a change in descending modulation will be evident in any area remote to the injured or painful site features a static nature in a rather dynamic phenomenon.<sup>36</sup> This raises questions regarding the choice of the most appropriate testing location which would be anatomically close to—yet sufficiently remote from—the painful region. The design of the present study did not allow for a comparison of the CPM at the painful versus a pain-free site. Comparison of the CPM effect as elicited from affected versus unaffected tissues would have a great potential in further enhancing our understanding of tendon pain pathophysiology and warrants future research.

### Is weaker conditioned pain modulation 'dysfunctional'?

Another point that merits attention is the label 'dysfunctional' that is given to CPM when found to be different between patient and healthy groups. Rationalising CPM teleologically by the speculative but widely accepted notion that pain acts as a warning system for danger or threat, the reduced CPM response pattern seen in patient populations might reflect an adaptive mechanism with the purpose to protect the painful area. For

such an adaptation to be functional, the assumption that a threat is implied to the body is required. It is possible that the reduced CPM effect in the Achilles tendinopathy group, as observed in this study, is the expression of a dynamic and flexible pain modulation process in the presence of increased afferent nociceptive input warning for tissue damage. However, this view is confronted by the common disassociation of tissue damage and pain.<sup>64–67</sup>

### CONCLUSION

Although the pain in tendinopathy is generally considered to be driven by peripheral mechanisms, the findings of this study reveal that people with Achilles tendinopathy modulate pain centrally in a different way than healthy people. Such differences are evident in a variety of chronic pain conditions and have been suggested to contribute to the chronicity of pain. Thus, the reduced CPM as observed in people with Achilles tendinopathy may promote ongoing tendon pain.

#### What are the findings?

- ▶ Pressure pain threshold testing over the Achilles tendon revealed that the conditioned pain modulation effect is reduced in people with Achilles tendinopathy.
- ▶ The findings indicate that besides peripheral mechanisms, altered central pain processing also plays a role in persistent Achilles tendinopathy.
- ▶ The reduction in conditioned pain modulation in people with Achilles tendinopathy was not correlated with the severity of Achilles tendinopathy (VISA-A), functional limitations (FAAM), and activity levels (amount of running).

#### How might it impact on clinical practice in the future?

- ▶ If altered central pain modulation in people with Achilles tendinopathy can be substantiated in future studies, the clinical examination for people with Achilles tendinopathy might have to incorporate assessment techniques to identify all relevant contributing pain mechanisms.
- ▶ If future research confirms altered central pain modulation in people with Achilles tendinopathy, it would be logical to explore drug therapy and treatment modalities that target the central nervous system in people with Achilles tendinopathy.
- ▶ Considering the presence of altered conditioned pain modulation effects in people with Achilles tendinopathy, similar findings in patients with chronic and widespread pain conditions, such as fibromyalgia and chronic fatigue syndrome, may need to be reinterpreted.

**Twitter** Follow Michel Coppieters at @michelcoppie

**Contributors** NT was responsible for the acquisition and analysis of the data, and drafting of the work. All authors contributed substantially to the conception and design of the work, and the interpretation of the data. All authors revised the work critically for intellectual integrity and provided final approval for the version to be published.

**Competing interests** None declared.

**Ethics approval** Ethics committee of the Department of Human Movement Sciences of VU University Amsterdam, The Netherlands (Reference: ECB 2014-58).

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** Raw data can be shared for research purposes on request.

## REFERENCES

- Maffulli N, Khan KM, Puddu G. Overuse tendon conditions: time to change a confusing terminology. *Arthroscopy* 1998;14:840–3.
- Järvinen TA, Kannus P, Maffulli N, et al. Achilles tendon disorders: etiology and epidemiology. *Foot Ankle Clin* 2005;10:255–66.
- Paavola M, Kannus P, Paakkala T, et al. Long-term prognosis of patients with Achilles tendinopathy. An observational 8-year follow-up study. *Am J Sports Med* 2000;28:634–42.
- Rio E, Kidgell D, Purdam C, et al. Isometric exercise induces analgesia and reduces inhibition in patellar tendinopathy. *Br J Sports Med* 2015;49:1277–83.
- Rio E, Moseley L, Purdam C, et al. The pain of tendinopathy: physiological or pathophysiological? *Sports Med* 2014;44:9–23.
- Heales LJ, Lim EC, Hodges PW, et al. Sensory and motor deficits exist on the non-injured side of patients with unilateral tendon pain and disability—implications for central nervous system involvement: a systematic review with meta-analysis. *Br J Sports Med* 2014;48:1400–6.
- Alfredson H, Spang C, Forsgren S. Unilateral surgical treatment for patients with midportion Achilles tendinopathy may result in bilateral recovery. *Br J Sports Med* 2014;48:1421–4.
- Coombes BK, Bisset L, Vicenzino B. A new integrative model of lateral epicondylalgia. *Br J Sports Med* 2009;43:252–8.
- Woolf CJ. Central sensitization: implications for the diagnosis and treatment of pain. *Pain* 2011;152:52–15.
- Staud R. Evidence for shared pain mechanisms in osteoarthritis, low back pain, and fibromyalgia. *Curr Rheumatol Rep* 2011;13:513–20.
- Van Oosterwijk J, Nijs J, Meeus M, et al. Evidence for central sensitization in chronic whiplash: a systematic literature review. *Eur J Pain* 2013;17:299–312.
- Skou ST, Graven-Nielsen T, Rasmussen S, et al. Widespread sensitization in patients with chronic pain after revision total knee arthroplasty. *Pain* 2013;154:1588–94.
- Staud R. Abnormal endogenous pain modulation is a shared characteristic of many chronic pain conditions. *Expert Rev Neurother* 2012;12:577–85.
- Fernandez-Carnero J, Fernandez-de-Las-Penas C, de la Llave-Rincon AI, et al. Widespread mechanical pain hypersensitivity as sign of central sensitization in unilateral epicondylalgia: a blinded, controlled study. *Clin J Pain* 2009;25:555–61.
- Maffulli N, Kenward MG, Testa V, et al. Clinical diagnosis of Achilles tendinopathy with tendinosis. *Clin J Sport Med* 2003;13:11–15.
- van Wilgen C, Konopka K, Keizer D, et al. Do patients with chronic patellar tendinopathy have an altered somatosensory profile? A Quantitative Sensory Testing (QST) study. *Scand J Med Sci Sports* 2013;23:149–55.
- Skinner IW, Debenham JR, Krumenachera S, et al. Chronic mid portion Achilles tendinopathy is not associated with central sensitisation. *Pain Rehabil - J Physiother Pain Assoc* 2014;2014:34–40.
- Kountouris A, Cook J. Rehabilitation of Achilles and patellar tendinopathies. *Best Pract Res Clin Rheumatol* 2007;21:295–316.
- Lewis GN, Heales L, Rice DA, et al. Reliability of the conditioned pain modulation paradigm to assess endogenous inhibitory pain pathways. *Pain Res Manag* 2012;17:98–102.
- Pud D, Granovsky Y, Yarnitsky D. The methodology of experimentally induced diffuse noxious inhibitory control (DNIC)-like effect in humans. *Pain* 2009;144:16–19.
- Tracey I, Mantyh PW. The cerebral signature for pain perception and its modulation. *Neuron* 2007;55:377–91.
- Julien N, Goffaux P, Arsenault P, et al. Widespread pain in fibromyalgia is related to a deficit of endogenous pain inhibition. *Pain* 2005;114:295–302.
- King CD, Wong F, Currie T, et al. Deficiency in endogenous modulation of prolonged heat pain in patients with irritable bowel syndrome and temporomandibular disorder. *Pain* 2009;143:172–8.
- Olesen SS, Brock C, Krarup AL, et al. Descending inhibitory pain modulation is impaired in patients with chronic pancreatitis. *Clin Gastroenterol Hepatol* 2010;8:724–30.
- Drummond PD, Knudsen L. Central pain modulation and scalp tenderness in frequent episodic tension-type headache. *Headache* 2011;51:375–83.
- Arendt-Nielsen L, Nie H, Laursen MB, et al. Sensitization in patients with painful knee osteoarthritis. *Pain* 2010;149:573–81.
- Corrêa JB, Costa LO, de Oliveira NT, et al. Central sensitization and changes in conditioned pain modulation in people with chronic nonspecific low back pain: a case-control study. *Exp Brain Res* 2015;233:2391–9.
- Hutchison AM, Evans R, Bodger O, et al. What is the best clinical test for Achilles tendinopathy? *Foot Ankle Surg* 2013;19:112–17.
- Robinson JM, Cook JL, Purdam C, et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med* 2001;35:335–41.
- Martin RL, Irrgang JJ, Burdett RG, et al. Evidence of validity for the Foot and Ankle Ability Measure (FAAM). *Foot Ankle Int* 2005;26:968–83.
- Mitchell LA, MacDonald RA, Brodie EE. Temperature and the cold pressor test. *J Pain* 2004;5:233–7.
- Kregel J, van Wilgen CP, Zwerver J. Pain assessment in patellar tendinopathy using pain pressure threshold algometry: an observational study. *Pain Medicine* 2013;14:1769–75.
- Yarnitsky D, Arendt-Nielsen L, Bouhassira D, et al. Recommendations on terminology and practice of psychophysical DNIC testing. *Eur J Pain* 2010;14:339.
- Lewis GN, Rice DA, McNair PJ. Conditioned pain modulation in populations with chronic pain: a systematic review and meta-analysis. *J Pain* 2012;13:936–44.
- van Wijk G, Veldhuijzen DS. Perspective on diffuse noxious inhibitory controls as a model of endogenous pain modulation in clinical pain syndromes. *J Pain* 2010;11:408–19.
- Yarnitsky D. Conditioned pain modulation (the diffuse noxious inhibitory control-like effect): its relevance for acute and chronic pain states. *Curr Opin Anaesthesiol* 2010;23:611–15.
- Normand E, Potvin S, Gaumond I, et al. Pain inhibition is deficient in chronic widespread pain but normal in major depressive disorder. *J Clin Psychiatry* 2011;72:219.
- Heymen S, Maixner W, Whitehead WE, et al. Central processing of noxious somatic stimuli in patients with irritable bowel syndrome compared with healthy controls. *Clin J Pain* 2010;26:104–9.
- Wilder-Smith CH, Robert-Yap J. Abnormal endogenous pain modulation and somatic and visceral hypersensitivity in female patients with irritable bowel syndrome. *World J Gastroenterol* 2007;13:3699–704.
- Oono Y, Wang K, Baad-Hansen L, et al. Conditioned pain modulation in temporomandibular disorders (TMD) pain patients. *Exp Brain Res* 2014;232:3111–19.
- Ng TS, Pedler A, Vicenzino B, et al. Less efficacious conditioned pain modulation and sensory hypersensitivity in chronic whiplash-associated disorders in Singapore. *Clin J Pain* 2014;30:436–42.
- Daenen L, Nijs J, Roussel N, et al. Dysfunctional pain inhibition in patients with chronic whiplash-associated disorders: an experimental study. *Clin Rheumatol* 2013;32:23–31.
- Cathcart S, Winefield AH, Lushington K, et al. Noxious inhibition of temporal summation is impaired in chronic tension-type headache. *Headache* 2010;50:403–12.
- Pielstick A, Haag G, Zaudig M, et al. Impairment of pain inhibition in chronic tension-type headache. *Pain* 2005;118:215–23.
- Kosek E, Ordeberg G. Lack of pressure pain modulation by heterotopic noxious conditioning stimulation in patients with painful osteoarthritis before, but not following, surgical pain relief. *Pain* 2000;88:69–78.
- Heinrich M, Tavares I, Leith J, et al. Descending control of nociception: specificity, recruitment and plasticity. *Brain Res Rev* 2009;60:214–25.
- Meeus M, Nijs J, Van de Wauwer N, et al. Diffuse noxious inhibitory control is delayed in chronic fatigue syndrome: an experimental study. *Pain* 2008;139:439–48.
- Vanegas H, Schaible H. Descending control of persistent pain: inhibitory or facilitatory? *Brain Res Brain Res Rev* 2004;46:295–309.
- Miranda J, Lamana S, Dias E, et al. Effect of pain chronification and chronic pain on an endogenous pain modulation circuit in rats. *Neuroscience* 2015;286:37–44.
- Yarnitsky D, Crispel Y, Eisenberg E, et al. Prediction of chronic post-operative pain: pre-operative DNIC testing identifies patients at risk. *Pain* 2008;138:22–8.
- Gwilym SE, Keltner JR, Warnaby CE, et al. Psychophysical and functional imaging evidence supporting the presence of central sensitization in a cohort of osteoarthritis patients. *Arthritis Rheum* 2009;61:1226–34.
- Fernández-de-Las-Peñas C, Galán-del-Río F, Fernández-Carnero J, et al. Bilateral widespread mechanical pain sensitivity in women with myofascial temporomandibular disorder: evidence of impairment in central nociceptive processing. *J Pain* 2009;10:1170–8.
- Price DD, Zhou Q, Moshiree B, et al. Peripheral and central contributions to hyperalgesia in irritable bowel syndrome. *J Pain* 2006;7:529–35.
- Vierck CJ. Mechanisms underlying development of spatially distributed chronic pain (fibromyalgia). *Pain* 2006;124:242–63.
- Giesecke T, Gracely RH, Grant MA, et al. Evidence of augmented central pain processing in idiopathic chronic low back pain. *Arthritis Rheum* 2004;50:613–23.
- Desmeules J, Cedraschi C, Rapiet E, et al. Neurophysiologic evidence for a central sensitization in patients with fibromyalgia. *Arthritis Rheum* 2003;48:1420–9.
- Curatolo M, Petersen-Felix S, Arendt-Nielsen L, et al. Central hypersensitivity in chronic pain after whiplash injury. *Clin J Pain* 2001;17:306–15.
- Bendtsen L. Central sensitization in tension-type headache—possible pathophysiological mechanisms. *Cephalalgia* 2000;20:486–508.
- Edwards RR, Ness TJ, Weigent DA, et al. Individual differences in diffuse noxious inhibitory controls (DNIC): association with clinical variables. *Pain* 2003;106:427–37.

## Original article

- 60 Nir RR, Yarnitsky D. Conditioned pain modulation. *Curr Opin Support Palliat Care* 2015;9:131–7.
- 61 Ge HY, Nie H, Graven-Nielsen T, *et al.* Descending pain modulation and its interaction with peripheral sensitization following sustained isometric muscle contraction in fibromyalgia. *Eur J Pain* 2012;16:196–203.
- 62 Graven-Nielsen T, Wodehouse T, Langford R, *et al.* Normalization of widespread hyperesthesia and facilitated spatial summation of deep-tissue pain in knee osteoarthritis patients after knee replacement. *Arthritis Rheum* 2012;64:2907–16.
- 63 Cook JL, Purdam CR. Is tendon pathology a continuum? A pathology model to explain the clinical presentation of load-induced tendinopathy. *Br J Sports Med* 2009;43:409–16.
- 64 Giombini A, Dragoni S, Di Cesare A, *et al.* Asymptomatic Achilles, patellar, and quadriceps tendinopathy: a longitudinal clinical and ultrasonographic study in elite fencers. *Scand J Med Sci Sports* 2013;23:311–16.
- 65 Comin J, Cook JL, Malliaras P, *et al.* The prevalence and clinical significance of sonographic tendon abnormalities in asymptomatic ballet dancers: a 24-month longitudinal study. *Br J Sports Med* 2013;47:89–92.
- 66 de Vos RJ, Heijboer MP, Weinans H, *et al.* Tendon structure's lack of relation to clinical outcome after eccentric exercises in chronic midportion Achilles tendinopathy. *J Sport Rehabil* 2012;21:34–43.
- 67 Malliaras P, Cook J. Patellar tendons with normal imaging and pain: change in imaging and pain status over a volleyball season. *Clin J Sport Med* 2006;16:388–91.



## Central pain processing is altered in people with Achilles tendinopathy

Nefeli Tompra, Jaap H van Dieën and Michel W Coppieters

*Br J Sports Med* published online December 23, 2015

---

Updated information and services can be found at:

<http://bjsm.bmj.com/content/early/2015/12/20/bjsports-2015-095476>

---

*These include:*

### References

This article cites 67 articles, 11 of which you can access for free at:  
<http://bjsm.bmj.com/content/early/2015/12/20/bjsports-2015-095476#BIBL>

### Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Topic Collections

Articles on similar topics can be found in the following collections

[Achilles tendinitis](#) (64)

---

### Notes

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>